PhytoKeys 75: 19–29 (2016)
doi: 10.3897/phytokeys.75.10244
http://phytokeys.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
PhytoKeys
Launched to accelerate biodiversity research

Galium shinasii (Rubiaceae): a new species of *Galium* L. from Eastern Turkey

Levent Şık[1], Hasan Yıldırım[2], Ademi Fahri Pirhan[2], Yusuf Altıoğlu[2], Meliha Gemici[2]

1 *Department of Biology, Faculty of Science and Arts, Celal Bayar University, 45030 Manisa, Turkey* **2** *Ege University, Faculty of Science, Department of Biology, 35100, Bornova-Izmir, Turkey*

Corresponding author: *Hasan Yıldırım* (hasanyldrm@gmail.com)

Academic editor: *S. Knapp* | Received 20 August 2016 | Accepted 21 November 2016 | Published 29 November 2016

Citation: Şık L, Yıldırım H, Pirhan AF, Altıoğlu Y, Gemici M (2016) *Galium shinasii* (Rubiaceae): a new species of *Galium* L. from Eastern Turkey. PhytoKeys 75: 19–29. doi: 10.3897/phytokeys.75.10244

Abstract

Galium shinasii Yıldırım (Rubiaceae), is described as a new species from Malatya Province in eastern Anatolia, Turkey. The new species is morphologically related to *Galium cornigerum* Boiss. & Hausskn. *G. lasiocarpum* and *G. sorgereae* Ehrend. and Schönb. but clearly differs from them based on the morphological differences presented in the species description. In addition, the conservation status, the distribution map, and notes on the biogeography and ecology of the new species are given.

Keywords

Rubiaceae, *Galium*, ecology, taxonomy, Turkey

Introduction

The family Rubiaceae is the fourth-biggest angiosperm family with about 660 genera and 11.500 species (Robbrecht and Manen 2006, Soza and Olmstead 2010). A large number of species of Rubiaceae are herbaceous and many are adapted to xeric habitats. (Robbrecht 1988, Jansen et al. 2000). *Galium* L. is one of the largest genera of the

Copyright *Levent Şık et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Rubiaceae, with about 650 species and approximately 780 taxa placed in 16 sections, including perennial and annual herbs that are distributed in temperate and tropical regions of worldwide (Willis 1985, Mabberley 1987, Goavert 2015).

A total of 121 *Galium* taxa (104 species) are found in Turkey and 60 taxa (endemism rate of 50%) are endemic to this country (Karabacak 2012).

Malatya is located in the eastern part of Turkey, one of the richest centres of species endemism in Turkey (Yıldız et al. 2004) and in recent years several new species have been described from this area and still continue to be discovered in Malatya (Yıldırım et al. 2010, Mutlu and Karakuş 2012, Tan et al. 2012, Koç and Aksoy 2013, Yıldırım and Erol 2013, Yıldırım and Şenol 2014a, 2014b, 2014c, Mutlu and Karakuş 2015a, 2015b, Yıldırım 2015a, 2015b).

Levent Canyon is one of the famous areas among biologists because of the high endemism in Malatya province. Recntly, several new plant species were described in this area. Levent Canyon is characterized by marlstone, a soft, finely fissured sedimentary rock (Schnurrenberger et al. 2003, Yıldırım and Şenol 2014c) which hosts many chasmophytes. The Levent Canyon is also a centre of diversity and endemism for several chasmophyte species. For example, *Reseda malatyana* Yıldırım (Resedaceae) and Şenol, *Alkanna malatyana* Şenol and Yıldırım (Boraginaceae), *Campanula alisan-kilincii* Yıldırım and Şenol (Campanulaceae), *Galium scopulorum* Schönb.-Tem. (Rubiaceae), *Pimpinella paucidentata* V.A.Matthews (Apiaceae), *Parietaria semispeluncaria* Yıldırım (Urticaceae) and *Galium cornigerum* Boiss. and Hausskn (Rubiaceae).

In June 2011, the second author collected an unusual and distinct specimen of *Galium* on marlstone-calcareous rocky cliffs in the Levent Canyon which authors believe to be of a new species for science.

Materials and methods

Specimens of the putative new species were compared with herbarium specimens at ANK, EGE, E, G, GAZI, HUB, ISTE, K, W and WU. In addition, the relevant literature (Halácsy 1901, Ehrendorfer and Krendl 1976, Ehrendorfer and Schönbeck-Temesy 1982, Davis et al. 1988, Ehrendorfer and Schönbeck-Temesy 2005, Govaerts 2016) was reviewed. The new species was examined by stereo-binocular microscope for morphological characterisation. At least 20 mature seeds and 30 pollen grains were measured using a light microscopy. For scanning electron microscopy (SEM), the selected seed and pollen grains were placed on aluminum stubs using double-sided adhesive tape, sputter coated with gold using an Emiteck K550, and then examined using the FEI Quanta250 FEG SEM. Photographs of living material were taken with a Nikon D300 digital camera. The conservation status of new species was evaluated based on the field observations in accordance with IUCN guidelines (2012). Geographical positions were recorded using a Magellan explorist 500 GPS.

Results

***Galium shinasii* Yıldırım, sp. nov.**
urn:lsid:ipni.org:names:77158796-1
Figure 1–3

Type. TURKEY: B7 Malatya: Akçadağ district, Levent Canyon, on marlstone rocky cliffs 1390 m, 26.06.2011, *H.Yıldırım 2128* (holotype: EGE42431!, isotypes: EGE42432!, NGBB!, ANK!).

Paratype. TURKEY: Malatya: Akçadağ district, Levent Canyon, on marlstone rocky cliffs 1390 m, 29.06.2015, H.Yıldırım 3358 (EGE!) (Figure 2A); loc., ibid., 11.09.2015, H.Yıldırım 3713(EGE!); Doğanşehir, Eskiköy, Meletbaşı mezrası karşısı kalker kayalıklar, 1630–1800 m., 12.07.2014, H.Yıldırım 3033 (EGE!) (Figure 2B). Erzincan: Sivas-Refahiye yolu, Refahiye'ye 1–2 km kala, kalker kaya üzeri, 1528 m, 09.09.2015, H.Yıldırım 3694 (EGE!) (Figure 2C).

Diagnosis. *Galium shinasii* is related to *Galium cornigerum*, *G. lasiocarpum* and *G. sorgereae* but it differs from them in having very reduced flowers (not flowers relatively larger), 1.2–1.8 mm corolla diam (not 2–5 mm); yellowish-green to reddish-green and 0,5–1 mm long tepals (not white or pink and not at least 2 mm); dorsal and ventral surface of with densely transparent tubercles and lateral surface 0.2–0.4 mm spreading to patent hairy fruits (not tubercles absent and fruits wholly villous, hirsute or subtomentose).

Description. Dwarf, caespitose perennial plant with many headed rootstock, suffruticose at base. Stem 1.5–6 cm long, fragile, prostrate-ascending to erect, many branched at base, glabrous to slightly puberulent, sometimes slightly winged on nerves, upper internodes elongate to 5 mm. Leaves in whorls of mostly 4, rarely 6, linear-lanceolate to narrow elliptic, 2–8 × 0.6–1.3 mm, 1 veined, glabrous to slightly puberulent, revolute at margin. Inflorescense dicashium, mostly terminal and also axillary, 8 to 75 flowered per stem; bracteoles absent. Pedicel glabrous, 1.5–2.5 mm in flowers, 2–5 mm in fruit. Calyx absent. Corolla 4 merious, yellowish-green to reddish-green, 1.2–1.8 mm diam; usually conical or campanulate, rarely infundibular; tube very reduced; lobes 0.5–1 × 0.4–0.7 mm, glabrous, triangular to lanceolate, mucronate at apex and apex incurved on petal inner surface. Stamen 0.4-0.6 mm long; anther yellow. Ovary 0.4–0.5 mm diam, dorsal and ventral surface of with densely transparent tubercles, lateral surface 0.2–0.4 mm spreading to patent hairy. Fruit depressed subglobose in fleshy, 0.5–0.75 mm, dorsal and ventral surface of with densely transparent tubercles, lateral surface 0.2–0.4 mm spreading to patent hairy.

Etymology. This species is named in honour of retired Prof. Dr. Şinasi Yıldırımlı (Biology Dep. Hacettepe University, Turkey), who is an expert in Plant systematics and taxonomy. He described more than 100 new plant species for science in Turkey. The Turkish name of this species is given as "Levent İplikçiği", according to the guidelines of Menemen et al. (2013).



Figure 1. *Galium shinasii*: **A** habit **B** flower **C** fruit **D** leaves.

Additional specimens examined. -*G. cornigerum*: –Turkey: Malatya: Levent Kanyonu inişi, kalker kayalıklar, 30.05.2012, H.Yıldırım 2386 (EGE!); Darende, Engüzek yaylası, Akbabaçalı Dağı zirve, 2100 m, 30.05.2012. H.Yıldırım 2396 (EGE!) (Figure 4D, E).



Figure 2. *Galium shinasii* habits: **A–B** from type locality, Levent Canyon, Malatya (H.Yıldırım 3358) **C–D** from Refahiye, Erzincan (H.Yıldırım 3694) **E–F** from Doğanşehir, Malatya (H.Yıldırım 3033).

-*G. incanum* subsp. *pseudocornigerum*: –Turkey: Sivas: Gürün-Pınarbaşı arası, Pınarbaşı'na 68 km kala, yol kenarı kayalık alan, 1662 m, 29.06.2016, H.Yıldırım 3948 (EGE!) (Figure 4B, C).

-*G.lasiocarpum*: –Turkey: Elazığ: in Cappadocia, Aucher 694 (holotype G!).

G. sorgerae: –Turkey: Isparta: Dedegöl Da., 2200 m, 1 vii 1965, Sorger 65-42- 70 (isotype WU!).

Distribution and ecology. *Galium shinasii* is a endemic for Eastern Anatolia. It's known that is from Levent Canyon (Figure 5) in Akçadağ district, and Eskiköy in



Figure 3. *Galium shinasii*: **A–C** flowers **D–E** fruits **F** fruit tubercles **G–J** pollen grains **K–M** detail of pollen garin surface.

Doğanşehir district in Malatya province and near Refahiye district in Erzincan (Figure 5). It is an element belonging to the Irano-Turanian floristic region and colonizes only marlstone-calcareous cliffs, usually those with an eastern and south-eastern orientation, at an elevation of 1200–1800 m. It is an obligate chasmophyte.

Suggested conservational status. The new species found in 3 populations. Two populations have been discovered in Malatya province, the other has been discovered in Erzincan province. Although the distribution area of *Galium shinasii* seems relatively wide, the populations of the area are very restricted. The total population area of *G. shinasii* was calculated as 0.2 km^2 and approximately 500 individuals were observed in total. Probably it has still several undiscovered populations. No anthropogenic or grazing effects were observed on the population. According to the present data, following the criteria laid out by the IUCN (2012), the plant is categorized as 'Vulnerable' (VU) D1 + 2, on account of its restricted distribution.

Discussion and conclusions

Galium shinasii is a member of *Galium* Sect. *Orientigalium* Ehrend. It is characterized by chasmophyte, dwarf caespitose habit, very fragile 2–6 cm long stems, mostly 4 rarely 6 leaves in a whorl; 1.5–6.5 cm long leaves; flowers diam 1.2–1.8 mm; corolla yellowish-green to reddish-green; usually corolla lobes formed in a conical or campanulate corolla shape, very rarely lobes wholly opens and formed a infundibular corolla shape; fruit dorsal and ventral surface with densely transparent tubercles, lateral surface 0.2–0.4 mm spreading to patent hairy.

Although *Galium shinasii* shows some morphological similarities with *G. lasiocarpum* Boiss., *G. sorgerae* Ehrend. and Schönb., *G. cornigerum* Boiss. and Hausk. in sect. *Orientigalium*, it is easily distinguished from these by relatively smaller flowers; yellowish-green to reddish-green and very reduced tepals; fruit surface is not only hairy on lateral surface, and also dorsal and ventral surface with densely transparent tubercles. Also it shows slight morphological similarities to *G. incanum* Sm. subsp. *pseudocornigerum* Ehrend. with dwarf caespitose habit, smaller leaves and in having fruits

Figure 4. A type locality of *Galium shinasii*, Marlstone rocky cliffs in Levent Canyon **B–C***G. incanum* subsp. *pseudocornigerum* **D–F** *G. cornigerum*

lacking a calyx but it is easily distinguished from *G. incanum*. subsp. *pseudocornigerum* by its especially more reduced and different coloured flowers, a greater numbers flower number per stem; smaller, depressed subglobose and long-hairy fruits.

Table 1. Main differantial characters among *Galium shinasii* and close related species *G. sorgerae*, *G. cornigerum* and *G. lasiocarpum*.

Species Characters	*Galium shinasii*	*G. sorgerae*	*G. cornigerum*	*G. lasiocarpum*
Stem	1.5–6 cm, prostrate-ascending to erect, glabrous to slightly puberulent; sometimes slightly winged on nerves,	3–4 cm, prostrate-ascending, densely hirsute	to 5 cm, prostrate-ascending, with very short, subvelutinous hairy	to 5 cm, erect to ascending, covered with straight spreading hair
Leaves	2–8 × 0.6–1.3 mm, linear-lanceolate to narrow elliptic; in whorls mostly 4, rarely 6 leafed	4–6 × 0.7–1 mm, linear-oblanceolate or narrowly elliptic; in whorls 6 leafed;	5–9 × 0.4–0.8 mm, linear, linear elliptic to lanceolate; in whorls 6 leafed	6–10 × 1–1.5 mm, linear-elliptic; in whorls 6 leafed
Inflorescence	very reduced, dicashium, mostly terminal and also axillar, 8 to 75 flowered per stem, never hidden by uppermost leaves	very reduced, few-flowered	mainly terminal reduced, corymbiform cymes, few-flowered, often ± hidden by uppermost leaves	very reduced, subumbellate-capitate, 3–8 flowered, hidden by uppermost leaves
Pedicel	1.5–2.5 mm, glabrous, 2–5 mm in fruiting time	2–3 mm, hirsute	0–4 mm, subvelutinous hairy	1–3 mm, hairy
Calyx	absent	–	2–4, subulate, persistent in fruit.	0–2
Corolla	yellowish-green to reddish-green; 1.2–1.8 mm diam;	whitish when dry, infundibular, 2–2.5 mm diam	white, 4–5 mm diam	white or pink, 3.5–4 mm diam
Fruit	dorsal and ventral surface with densely transparent tubercles, lateral surface 0.2–0.4 mm spreading to patent hairy.	hirsute	subtomentose	villous



Figure 5. Known distribution of: *Galium shinasii* (✹), *G. cornigerum* (▲), *G. sorgerae* (●), *G. lasiocarpum* (■).

The detailed of the morphological differences between *Galium shinasii* and related *Gallium* species are summarized in the Table 1.

After adding this new species in science literature, the total number of *Gallium* taxa were raised to 121 (105 species) in Turkey and 61 taxa are endemic for Turkey.

Acknowledgements

We are grateful to the curators of the following herbaria for allowing us to access to their *Gallium* material for study: ANK, EGE, E, G, GAZI, HUB, ISTE,K, W and WU. The authors are indebted to The Scientific and Technological Research Council of Turkey (TUBITAK), under Project number: 133Z072, for the financial support.

References

Davis PH, Mill RR, Tan K (1988) Flora of Turkey and the East Aegean Islands (Vol. 10, Suppl.). Edinburgh University Press, Edinburgh, 216 pp.

Ehrendorfer F, Krendl F (1976) *Galium* L. In: Tutin TG, Heywood VH, Burges NA, Valentine DH (Eds) Flora Europaea, Vol. 4: Plantaginaceae to Compositae (and Rubiaceae). Cambridge University Press, Cambridge, 14–36.

Ehrendorfer F, Schönbeck-Temesy E (1982) *Galium* L. In: Davis PH (Eds) Flora of Turkey, Volume 7: Flora of Turkey and the East Aegean Islands, Vol. 7. Edinburgh University Press, Edinburgh, 767–849.

Ehrendorfer F, Schönbeck-Temesy E (2005) *Galium* L. In: Ehrendorfer F, Schonbeck-Temesy E, Puff C, Rechinger W (Eds) Flora Iranica (Volume 176). Verlag des naturhistorischen Museum Wien, Wien, 161–258.

Halácsy EV (1901) Conspectus Florae Graecae (Vol. 1). Sumptibus Guilelmi Engelmann, Lipsiae (Leipzig), 720–721.

Govaerts R (2016) World Checklist of Asparagaceae. Facilitated by the Royal Botanic Gardens, Kew. http://apps.kew.org/wcsp [retrieved 15/08/2016]

Jansen S, Dessein S, Piesschaert F, Robbrecht E, Smets E (2000) Aluminum accumulation in leaves of Rubiaceae: Systematic and phylogenetic implications. Annals of Botany 85: 91–101. doi: 10.1006/anbo.1999.1000

Karabacak E (2012) *Galium* L. In: Güner A, Aslan S, Ekim T, Vural M, Babaç MT (Eds) Türkiye Bitkileri Listesi (Damarlı Bitkiler). Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, 821–831.

Koç M, Aksoy A (2013) *Minuartia hamzaoglui* (Caryophyllaceae), a new species from Turkey. Turkish Journal of Botany 37: 428–433.

Mabberley DJ (1987) The Plant-Book: A Portable Dictionary of the Higher Plants. Cambridge University Press, Cambridge, 720 pp.

Menemen Y, Aytaç Z, Kandemir A (2013) Türkçe Bilimsel Bitki Adları Yönergesi, Bağbahçe Bilim Dergisi 47: 28–31.

Mutlu B, Karakuş Ş (2012) A new species of *Ornithogalum* (Hyacinthaceae) from East Anatolia, Turkey. Turkish Journal of Botany 36: 125–133.

Mutlu B, Karakuş Ş (2015a) A new species of *Campanula* (Campanulaceae) from Turkey. Phytotaxa 234(3): 287–293.

Mutlu B, Karakuş Ş (2015b) A new species of *Sisymbrium* (Brassicaceae) from Turkey: Morphological and molecular evidences. Turkish Journal of Botany 39: 325–333

Robbrecht E (1988) Tropical Woody Rubiaceae – Characteristic Features and Progressions – Contributions to a New Subfamilial Classification. Volume 1, Opera Botanica Belgica, 271 pp.

Robbrecht E, Manen JF (2006) The major evolutionary lineages of the coffee family (Rubiaceae, angiosperms). Combined analysis (nDNA and cpDNA) to infer the position of *Coptosapelta* and *Luculia*, and supertree construction based on rbcL, rps16, trnLtrnF and atpB-rbcL data. A new classification in two subfamilies Cinchonoideae and Rubioideae. Systematics and Geography of Plants 76: 85–146.

Schnurrenberger DW, Russell JM, Kelts KR (2003) Classification of lacustrine sediments based on sedimentary components. Journal of Paleolimnology 29: 141–154. doi: 10.1023/A:1023270324800

Soza VL, Olmstead RG (2010) Evolution breeding systems and fruits in New World *Galium* and relatives (Rubiaceae). American Journal of Botany 97: 1630–1646. doi: 10.3732/ajb.1000130

Tan K, Yıldırım H, Zielinski J (2012) Establishment of *Rosa* sect. *Caninae* subsect. *Orientales* (Rosaceae) and the recognition of an unusual variety of *Rosa vanheurckiana* from eastern Anatolia, Turkey. Phytotaxa 54: 26–36. doi: 10.11646/phytotaxa.54.1.3

Willis JC (1985) A Dictionary of the Flowering Plants and Ferns. Cambridge University Press, Cambridge, 1294 pp.

Yıldırım H, Tan K, Şenol SG, Pirhan AF (2010) *Chaenorhinum semispeluncarum* sp. nov. and *C. yildirimlii* sp. nov. (Scrophulariaceae) from east Anatolia, Turkey. Nordic Journal of Botany 28: 457–464. doi: 10.1111/j.1756-1051.2010.00790.x

Yıldırım H, Erol O (2013) *Crocus yakarianus* sp. nov. from eastern Turkey. Nordic Journal of Botany 31: 426–429. doi: 10.1111/j.1756-1051.2012.01786.x

Yıldırım H, Şenol SG (2014a) *Campanula alisan-kilincii* (Campanulaceae), a new species from eastern Anatolia, Turkey. Turkish Journal of Botany 38: 22–30.

Yıldırım H, Şenol SG (2014b) *Alkanna malatyana* (Boraginaceae), a new species from East Anatolia, Turkey. Phytotaxa 164(2): 124–132.

Yıldırım H, Şenol SG (2014c) *Reseda malatyana* (Resedaceae), a new chasmophytic species from eastern Anatolia, Turkey. Turkish Journal of Botany 38: 1013–1021.

Yıldırım H (2015a) *Muscari atillae* (Asparagaceae): a new species from Eastern Anatolia, Turkey. Pyhtotaxa 213(3): 291–295.

Yıldırım H (2015b) *Parietaria semispeluncaria* (Urticaceae): a new species from Eastern Anatolia, Turkey. Pyhtotaxa 226(3): 281–287.

Yıldız B, Bahçecioğlu Z, Arabacı T (2004) Floristic characteristics of Beydağı (Malatya). Turkish Journal of Botany 28: 391–419.